

# Square lattice site percolation at increasing ranges of neighbor interactions

Krzysztof Malarz<sup>1,\*</sup> and Serge Galam<sup>2,†</sup>

<sup>1</sup> AGH University of Science and Technology, Faculty of Physics and Applied Computer Science  
al. Mickiewicza 30, PL-30059 Kraków, Poland

<sup>2</sup> Université Pierre et Marie Curie et CNRS, Laboratoire des Milieux Désordonnés et Hétérogènes  
Case 86, 4 place Jussieu, F-75252 Paris Cedex 05, France

(Dated: February 2, 2008)

We report site percolation thresholds for square lattice with neighbor interactions at various increasing ranges. Using Monte Carlo techniques we found that nearest neighbors ( $N^2$ ), next nearest neighbors ( $N^3$ ), next next nearest neighbors ( $N^4$ ) and fifth nearest neighbors ( $N^6$ ) yield the same  $p_c = 0.592 \dots$ . At odds, fourth nearest neighbors ( $N^5$ ) give  $p_c = 0.298 \dots$ . These results are given an explanation in terms of symmetry arguments. We then consider combinations of various ranges of interactions with ( $N^2+N^3$ ), ( $N^2+N^4$ ), ( $N^2+N^3+N^4$ ) and ( $N^2+N^5$ ). The calculated associated thresholds are respectively  $p_c = 0.407 \dots$ ,  $0.337 \dots$ ,  $0.288 \dots$ ,  $0.234 \dots$ . The existing Galam–Mauger universal formula for percolation thresholds does not reproduce the data showing dimension and coordination number are not sufficient to build a universal law which extends to complex lattices.

PACS numbers: 07.05.Tp, 64.60.Ak

## I. INTRODUCTION

Calculating percolation thresholds has been an ongoing challenge for decades [1, 2, 3, 4, 5, 6]. While very few lattices allow an exact analytical calculation, large scale simulations have been very valuable to determine a large spectrum of them for both Bravais' [4, 7, 8] and disordered [9] lattices. The drastic increase in computer capacities has recently permitted the calculation of thresholds at rather high dimensions up to  $d = 13$  for the hypercube [4, 10]. In parallel, not much work has been devoted to regular lattices with neighbor interactions which are not nearest neighbors ( $N^2$ , von Neumann's neighborhood). Some scarce results are available for simultaneous nearest and next nearest neighbors ( $N^2+N^3$ , Moore's neighborhood) [4, 11].

In this paper we report for the first time a systematic calculation of site percolation thresholds for the square lattice with neighbor interactions at successive increasing range. We consider the series of nearest neighbors ( $N^2$ ), next nearest neighbors ( $N^3$ ), next next nearest neighbors ( $N^4$ ), fourth nearest neighbors ( $N^5$ ) and fifth nearest neighbors ( $N^6$ ). It should be stressed that for each one of the considered distance of interaction, all others are not active. For instance in the case of next nearest neighbors ( $N^3$ ), the nearest neighbors ( $N^2$ ) sites are not connected, only the  $N^3$  are. This principle applies to all our calculations. We found that the threshold is the same for all of them with  $p_c = p_c(N^2)$  except at  $N^5$ . An explanation in terms of symmetry is provided.

We then consider combinations of various ranges of interactions with ( $N^2+N^3$ ), ( $N^2+N^4$ ), ( $N^2+N^5$ ) and ( $N^2+N^3+N^4$ ). In these cases we have simultaneous range

of interactions but they are necessary compact. For instance for ( $N^2+N^4$ ) all nearest neighbors sites are connected as well as all next next nearest neighbor ones but next nearest neighbors are not interacting.

Comparing our numerical estimates with the predictions from the Galam–Mauger (GM) universal formula for percolation thresholds [12], we found significant discrepancies. It strengthens the earlier claim that only dimension and coordination number could not be sufficient to build a universal law which extends to complex lattices [7, 13].

## II. CALCULATIONS

There exist several computational techniques which allow to perform calculations of percolation thresholds [14, 15, 16, 17]. Here we are using the Hoshen–Kopelman algorithm (HKA) [17]. Once the lattice is given with the occupied sites, it allows to recognize which sites belong to which clusters. With HKA one can assign to each occupied site a label and sites in the same cluster have the same labels. Different labels are assigned to different clusters. The HKA is particularly efficient when we check if the site at distance  $\ell$  from the first line — often fully occupied — is still connected to that line through the sites at the distances smaller than  $\ell$ . The algorithm requires storing only single line of sites and goes through the lattice only once. In such a case HKA becomes extremely efficient as it saves memory and time [4]. However, when links between sites at distances larger than  $\ell$  from a top border are desired, the whole lattice must be stored [4, 18, 19, 20]. With the HKA on a square lattice when we assign the labels for the investigated site (black sites in Fig. 1), we need to check already labeled and occupied sites in its neighborhood (slashed sites in Fig. 1). The possible links to remaining sites in the neighborhood (backslashed sites in Fig. 1) may be checked later,

\*URL: <http://home.agh.edu.pl/malarz/>

†Electronic address: galam@ccr.jussieu.fr




FIG. 1: Various site neighborhoods on the square lattice: (a)  $N^2$  — von Neumann's neighborhood, (b)  $N^3$ , (c)  $N^4$ , (d)  $N^5$ , (e)  $N^6$ , and the examples of their combinations: (f)  $N^2+N^3$  — Moore's neighborhood, (g)  $N^2+N^4$ , (h)  $N^2+N^5$  and (i)  $N^2+N^3+N^4$ .

basing on the neighborhood's point symmetry.

The percolation thresholds values  $p_c$  are evaluated from the crossing point of three curves showing dependences of the percolation probability  $P$  on the site occupation probability  $p$  for lattices of linear sizes  $L = 100, 500$  and  $1000$ . The results are averaged over  $N_{\text{run}} = 10^3$  and  $10^4$  for  $L = 1000$  and  $100$ , respectively. With enlarging the lattice size  $L$  the curve  $P(p)$  becomes steeper and steeper and tends to Heaviside's function  $\Theta(-p_c)$  when  $L \rightarrow \infty$ , as expected.

### III. RESULTS

We present our results in Tab. I. The percolation thresholds  $p_c$  for the square lattice are computed with HKA for a series of neighborhoods. First only one type of neighbor interactions is considered at a time, increasing repeatedly the range with  $N^2$ ,  $N^3$ ,  $N^4$ ,  $N^5$  and  $N^6$ . It turns out that the threshold  $p_c = 0.592 \dots$  is the same for all of them except at  $N^5$  where  $p_c = 0.298 \dots$ .

Indeed, all lattices with neighborhoods shown in Figs. 1(b), (c) and (e) may be mapped into a  $N^2$  situation as in Fig. 1(a). The only difference is a larger and larger lattice constant. To implement the mapping, we take a square lattice and build on it the lattice from only  $N^3$  interactions. Two independent interpenetrated squares sublattices appear. Therefore the percolation of  $N^3$  is split onto two parallel  $N^2$  problems on each one of these two square sublattices. Accordingly the  $p_c$  on each one is the  $p_c$  of  $N^2$ . Moreover, as the site must be distributed homogeneously on the initial lattice, we will have the

TABLE I: The percolation threshold  $p_c$  for various neighborhoods on square lattice and sites coordination number  $z$  and the theoretical values  $p_c^{\text{GM}}$ .

neighborhood	$z$	$p_c$	$p_c^{\text{GM}}$
$N^2$	4	$0.592 \dots$	$0.5984 \dots$
$N^3$	4	$0.592 \dots$	$0.5984 \dots$
$N^4$	4	$0.592 \dots$	$0.5984 \dots$
$N^5$	8	$0.298 \dots$	$0.4411 \dots$
$N^6$	4	$0.592 \dots$	$0.5984 \dots$
$N^2+N^3$	8	$0.407 \dots$	$0.4411 \dots$
$N^2+N^4$	8	$0.337 \dots$	$0.4411 \dots$
$N^2+N^5$	12	$0.234 \dots$	$0.3748 \dots$
$N^2+N^3+N^4$	12	$0.288 \dots$	$0.3748 \dots$




FIG. 2: The lattices with (a)  $N^3$  and (b)  $N^4$  neighborhoods may be mapped to (a) two or (b) four parallel  $N^2$  situations but with (a)  $\sqrt{2}$  and (b) two times larger lattice constants.

same density of occupied sites on each one of the sublattice making both percolation to occur simultaneously at the same  $p_c$  (see Fig. 2(a)). Such a scheme can be repeated with  $N^4$  (Fig. 2(b)) and  $N^6$  but not with  $N^5$ . As shown in Fig. 1(d) the  $N^5$  lattice has eight neighbors while  $N^2$ ,  $N^3$ ,  $N^4$  and  $N^6$  have four.

These symmetry properties may become instrumental in underlining interesting physical properties associated to some exotic materials. In particular if one is able to discriminate between the two interpenetrated lattices, it may open a way to reach percolation at a much lower critical density, down to half the value of  $p_c$ . But such a search is out the scope of the present work.

We also consider several combinations of various ranges of interactions. First an increasing compact neighborhood with  $(N^2+N^3)$  and  $(N^2+N^3+N^4)$ . The calculated threshold numerical estimates are respectively  $p_c = 0.407 \dots$  and  $0.288 \dots$  (Tab. I). Then more complex ones with  $(N^2+N^4)$  and  $(N^2+N^5)$  for which we obtained  $p_c = 0.337 \dots$  and  $0.234 \dots$  (Tab. I). The fact that  $p_c$  of  $(N^2+N^5)$  is smaller than  $p_c$  of  $(N^2+N^3+N^4)$  is consistent with  $N^5$   $z = 8$  instead of  $z = 4$  for all the others.

The obtained percolation threshold values  $p_c(N^2) =$

$0.592 \dots$  and  $p_c(N^2 + N^3) = 1 - p_c(N^2) = 0.407 \dots$  are consistent with the values reported in Refs. 4, 11, 21. We have also revised the value of  $p_c(N^2 + N^3 + N^4)$  which has been studied in an earlier paper [22] and it was put at 0.292 to compare to our value of  $0.288 \dots$ .

#### IV. DISCUSSION

At this stage it is interesting to check the validity GM universal formula for percolation thresholds [12, 23] in the case of these complex neighbor interactions. Comparing our numerical estimates with its predictions as shown in Tab. I we found a good agreement for  $N^2$ ,  $N^3$ ,  $N^4$  and  $N^6$  ( $\Delta = 0.006$ ) but not for  $N^5$  ( $\Delta = 0.123$ ). It is also fair for  $(N^2 + N^3)$  with  $\Delta = 0.034$  but not for all others combinations. The significant discrepancies occur for complex and non compact neighborhood. It strengthens the earlier claim that only dimension and coordination number could not be sufficient to build a universal law which extends to complex lattices [13].

Indeed above failures could be anticipated due to the fact that several lattices have both identical  $z$  and  $d$  though they exhibit different thresholds as seen from Tab. I. In particular  $N^5$ ,  $(N^2 + N^3)$ ,  $(N^2 + N^4)$  have  $z = 8$  and  $d = 2$  while all  $p_c$  are different. The same occurs for  $(N^2 + N^5)$  and  $(N^2 + N^3 + N^4)$  with  $z = 12$  and  $d = 2$ .

Similar situation occurs for  $T_C$  in the Ising model where even with the same number of interacting spins in the neighborhood and the same dimensionality we have different  $T_C$  [24]. On the other hand Bragg-Williams approximation [25] predicts  $T_C$  to be unique function of

coordination number  $z$ , i.e.  $k_B T_C = zJ$  [26]. The GM universal formula which also extends to  $T_C$  includes a dependence on both  $d$  and  $z$  [27].

To conclude we have report for the first time numerical estimates for site percolation thresholds for the square lattice with  $N^3$ ,  $N^4$ ,  $N^5$ ,  $N^6$ ,  $(N^2 + N^4)$  and  $(N^2 + N^5)$  interactions. Our new estimates may prove useful in the search for a robust universal formula for percolation thresholds which would apply to complex lattices. In particular on how to extend the GM law by including some additional topological ingredient besides coordination  $z$  and dimension  $d$ .

These results may prove useful to some of the large spectrum of physical and interdisciplinary topics where the percolation theory may be applied like forest fires spreading [20, 28], immunology [29], liquid migration in porous media [30], econophysics [31], and sociophysics [32].

#### Acknowledgments

We would like to thank UnProf. Dr. habil Dipl.-Phys. D. Stauffer for introducing us each other during the 18<sup>th</sup> Max Born Symposium in Łądek Zdrój, 2003. KM's stay at the Université Pierre et Marie Curie et CNRS, LMDH was financed by the European Science Foundation with grant No. COST-STSM-P10-00262. The numerical calculations were carried out in ACK-CYFRONET-AGH. The machine time on SGI 2800 is financed by the Polish Ministry of Science and Information Technology under grant No. MNiI/SGI2800/AGH/049/2003.

- 
- [1] M. F. Sykes, M. Glen, J. Phys. A **9**, 87 (1976); M. F. Sykes, D. S. Gaunt, M. Glen, *ibid.* **9**, 97 (1976); M. F. Sykes, D. S. Gaunt, M. Glen, *ibid.* **9**, 715 (1976); M. F. Sykes, D. S. Gaunt, M. Glen, *ibid.* **9**, 725 (1976); D. S. Gaunt, M. F. Sykes, *ibid.* **9**, 1109 (1976).
  - [2] H. Kesten, *Percolation Theory for Mathematicians*, (Birkhauser, Boston, 1982).
  - [3] D. Stauffer, *Percolation in Fractals and Disordered Systems*, Eds. A. Bunde and S. Hauling, (Springer, Berlin, 1991).
  - [4] D. Stauffer, A. Aharony, *Introduction to Percolation Theory*, (Taylor and Francis, London, 1994).
  - [5] M. Sahimi, *Applications of Percolation Theory*, (Taylor and Francis, London, 1994).
  - [6] D. Stauffer, *Physica A* **242**, 1 (1997).
  - [7] P. N. Sinding, R. M. Ziff, *Phys. Rev. E* **60**, 275 (1999).
  - [8] A. Rosowsky, *Eur. Phys. J. B* **15**, 77 (2000).
  - [9] M. E. J. Newman, I. Jensen, R. M. Ziff, *Phys. Rev. E* **65**, 021904 (2002); D. S. Callaway, M. E. J. Newman, S. H. Strogatz, D. J. Watts, *Phys. Rev. Lett.* **85**, 5468 (2000); C. Moore, M. E. J. Newman, *Phys. Rev. E* **62**, 5678 (2000); C. Moore, M. E. J. Newman, *ibid.* **62**, 7059 (2000).
  - [10] S. Galam, A. Mauger, *Physica A* **205**, 502 (1994); P. Grassberger, *Phys. Rev. E* **67**, 036101 (2003).
  - [11] I. V. Petrov, I. I. Stoynev, F. V. Babalievskii, *J. Phys. A* **24**, 4421 (1991).
  - [12] S. Galam, A. Mauger, *Phys. Rev. E* **53**, 2177 (1996); S. Galam, A. Mauger, *Phys. Rev. E* **56**, 322 (1997); S. Galam, A. Mauger, *Eur. Phys. J. B* **1**, 255 (1998).
  - [13] S. C. van der Marck, *Phys. Rev. E* **55**, 1228 (1997); S. C. van der Marck, *ibid.* **55**, 1514 (1997); S. C. van der Marck, *ibid.* **55**, 3732(E) (1997); S. C. van der Marck, *Int. J. Mod. Phys. C* **9**, 529 (1998); F. Babalivski, *Phys. Rev. E* **59**, 1278 (1999).
  - [14] M. E. J. Newman, R. M. Ziff, *Phys. Rev. E* **64**, 016706 (2001); *Phys. Rev. Lett.* **85**, 4104 (2000).
  - [15] R. M. Ziff, *J. Stat. Phys.* **28**, 838 (1982).
  - [16] P. L. Leath, *Phys. Rev. B* **14**, 5046 (1976).
  - [17] J. Hoshen, R. Kopelman, *Phys. Rev. B* **14**, 3428 (1976).
  - [18] K. Malarz, A. M. Vidales, *Int. J. Mod. Phys. C* **9**, 147 (1998).
  - [19] D. Stauffer, A. Aharony, *Int. J. Mod. Phys. C* **10**, 935 (1999).
  - [20] K. Malarz, S. Kaczanowska, K. Kułakowski, *Int. J. Mod. Phys. C* **13**, 1017 (2002).
  - [21] R. M. Ziff, *Phys. Rev. Lett.* **69**, 2670 (1992).
  - [22] C. Domb, N. W. Dalton, *Proc. Phys. Soc.* **89**, 859 (1966).

- [23] J. C. Wierman, Phys. Rev. E **66**, 27105 (2002).
- [24] K. Malarz, Int. J. Mod. Phys. C **13**, 1017 (2002).
- [25] W. L. Bragg, E. J. Williams, Proc. Roy. Soc. A **145**, 699 (1934).
- [26] K. Huang, *Statistical Mechanics*, (John Wiley and Sons, Inc., New York, 1963).
- [27] S. Galam, A. Mauger, Physica A **235**, 573 (1997).
- [28] G. Caldarelli, *at al.*, `cond-mat/0108011`; J. A. M. S. Duarte, Ann. Rev. Comput. Phys. **V**, 1 (1997).
- [29] A. Jimenez-Dalmaroni, H. Hinrichsen, `cond-mat/0304113`; S. M. Dammer, H. Hinrichsen, `cond-mat/0303467`; L. M. Sander, C. P. Warren, I. M. Sokolov, `cond-mat/0301394`.
- [30] Y. Lee *at al.*, Phys. Rev. E **60**, 3425 (1999); A. M. Viales *at al.*, Europhys. Lett. **36**, 259 (1996).
- [31] A. Barrañón, `nlin.A0/0404009`; M. Hohnisch, S. Pittnauer, D. Stauffer, `cond-mat/0308358`; D. Makowiec, P. Gnaciński, W. Miklaszeski, `cond-mat/0307290`; J. Goldenberg, B. Libai, S. Solomon, N. Jan, D. Stauffer, `cond-mat/0005426`.
- [32] T. Erez, S. Moldovan, S. Solomon, `cond-mat/0406695`; S. Galam, A. Mauger, Physica A **323**, 695 (2003); S. Galam, Physica A **336**, 49 (2004); S. Galam, Eur. Phys. J. B **26**, 269 (2002); D. Stauffer, `cond-mat/0204099`; S. Galam, `cond-mat/0204052`; A. Proykova, D. Stauffer, `cond-mat/0203375`; S. Solomon, G. Weisbuch, L. de Arcangelis, N. Jan, D. Stauffer, Physica A **277**, 239 (2000).